Studies and sources

Study 1

Soft-tissue reaction

- Ceramic abutments show healthy and <u>stable gingiva</u> and proved to be superior to other materials. Andersson B, Taylor A, Lang BR, et al; Int J Prosthodont. 2001 Sep-Oct.
- Fibroblasts show superior connection to rough ceramic surfaces leading to connective <u>tissue attachment</u>. *Mustafa K, Oden A, Wennerberg A, et al; Biomaterials. 2005 Feb.*
- Attachment complex of gingiva and abutments made of aluminium-oxid ceramic is similar to that to the tooth. Exhibits a basal lamina and hemidesmosomal layer. *McKinney RV Jr, Steflik DE, Koth DL; J Periodontol. 1985 Oct.*
- <u>Plaque adhesion</u> to ceramic abutments is significantly less than to titanium abutments. Barclay CW, Last KS, Williams R; Int J Prosthodont. 1996 Sep-Oct.
- Positive tissue reaction to zirconium-oxide ceramic. Glauser R, Sailer I, Wohlwend A, et al; Int J Prosthodont. 2004 May-Jun.

Study 2

The natural model

- The phenomenon of <u>self cleansing</u> has been scrutinized and offers possibilities of protection <u>against plaque and microorganisms</u>. Applying the characteristics of the lotus effect on implant surfaces creates a self-cleansing phenomenon. *Furstner R, Barthlott W, Neinhuis C, et al; Langmuir. 2005 Feb.*

Study 3

- Two-piece implants show significant potential for inflammation and bone resorption at the micro gap. One-piece implants <u>showed none</u>. Broggini N, McManus LM, Hermann JS, et al; J Dent Res. 2003 Mar.
- One-piece implant show the least bone resorption when the surface is roughened. Surface roughness is carried to the margin of the crown, respecting the <u>biological width</u>. *Hartman GA, Cochran DL; J Periodontol. 2004 Apr.*
- With the two-piece system, <u>it is impossible to keep the micro-gap free of bacteria</u>. This leads to an inflammation of the marginal mucosa. *Guindy JS, Besimo CE, Besimo R, et al; J Oral Rehabil. 1998 Jun.*
- Sealing Materials to fill up the inside of an implant e.g. wax offer best nutrition for bacteria in the micro gap *Xin Xie; Universität Köln. 2004 Mai.*
- Positive <u>gingival aesthetics is based on a constant and healthy biological width</u>. The gingiva around the one piece implant is similar to the gingiva condition of natural teeth. Two piece systems show inferior biological results.
- Hermann JS, Buser D, Schenk RK, et al; Clin Oral Implants Res. 2001 Dec.

Study to 4

- The <u>optimal primary stability</u> of an implant screw in the human bone without fracture is achieved at 70 Ncm insertion torque within bicortical bone and 50 Ncm within unicortical bone. *Ueda M, Matsuki M, Jacobsson M, et al; Int J Oral Maxillofac Implants. 1991 Winter.*
- A moderate implant loading with provisional restorations improves the <u>perioimplant bone quality</u>. *Nentwig G H, Psenicka H; Göttingen. 2003 Nov.*

Contact:

Dr. Peter Bongard Specialist (EDA, BDIZ) for Implantology Periodontology Länglingsweg 69 b, 47441 Moers, Germany fon: 0 28 41 - 3 10 40, fax: 0 28 41 - 3 32 72 e-Mail: praxisbongard@t-online.de

Does the perfect implant exist?


Result of scientific studies. The One Piece Ceramic-Titanium Implant. (patent.)



The One Piece Ceramic-Titanium Implant.

(patent.)

The biological, aesthetic implant that follows the model of nature.



due to the individual preparation of the ceramic in harmony with

Stable connection of the ceramic to the underlying titanium when exposed to

No horizontal and vertical bone-loss caused by microorganisms as found in the